

A cleavable chimeric peptide-R7: enhancing the dual-function of LPS neutralization and antibacterial properties against MDR E. coli

Zhenlong Wang^{1,2}, Da Teng^{1,2}, Ruoyu Mao^{1,2}, Ya Hao^{1,2}, Na Yang^{1,2}, Xiumin Wang^{1,2}, Jianhua Wang^{1,2}

¹ Key Laboratory of Feed Biotechnology, Ministry of Agriculture and Rural Affairs, Beijing 100081, People's Republic of China

² Gene Engineering Laboratory, Feed Research Institute, Chinese Academy of Agricultural Sciences, Beijing 100081, People's Republic of China

Copyright: 2021 Wang Z. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

During the last ten years, the design of new enzyme-free electrochemical sensors based metal oxide nanoparticles has been described. Due to the electrocatalytic activity of these nanomaterials, these sensors allow the sensitive detection of different ions and molecules. They can replace advantageously enzymes due to their higher stability.

The presentation will be focused on the use of Kegging type polyoxometallates (POMs) for the detection of nitrite ions¹ and of hydrogen peroxide. The association of POMs with single-wall carbon nanotubes (image) allowed the increase of the sensitivity of detection of hydrogen peroxide by a factor of 38.52. The involved charge transfer mechanism will be detailed.

Other types of metal oxide nanomaterials such as perovskites will also be presented. A₂B_{0.4}O_{4+δ} type perovskite are alternance of perovskite ABO₃ and of rock salt AO layers, A being a lanthanide and B a metal. Detection limit of glucose is 0.5 μM and no interference of detection is observed in human serum. The involved charge transfer mechanism will be detailed.

This presentation shows the potentiality of metal oxide nanomaterials for biosensing.

Biography:

Xiumin Wang has been focusing on this special narrow field of AMP for many years, especially design of different antibacterial agents, antibacterial activity, and mode of actions.

Citation: Xiumin Wang; A cleavable chimeric peptide-R7: enhancing the dual-function of LPS neutralization and antibacterial properties against MDR E. coli | International Webinar on Materials Science and Nanotechnology | February 25, 2021