

Efficacy of Ciclopirox Olamine cream in Superficial Dermatophytosis Affiliation: Dhaka Medical College hospital

Sharmin Jahan

Dhaka Medical College hospital, Bangladesh

Copyright: 2021 Sharmin J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Ciclopirox is a broad-spectrum antifungal agent as well as anti-inflammatory agent and is effective against both Gram-positive and Gram-negative bacteria. These properties may give ciclopirox an advantage over other antifungal agents in the treatment of dermatophyte infection.

Objective: To evaluate the efficacy and safety of Ciclopirox olamine 1% cream in the treatment of Dermatophytosis.

Methods: In Dhaka Medical College hospital Eighty-one patients with tinea corporis/tinea cruris were selected according to the inclusion and exclusion criteria and were examined for clinical diagnosis and confirmed by a positive 10.0% KOH reading. They were advised to apply Ciclopirox olamine 1% cream once daily for 2 weeks and Clinical and mycological assessment at baseline and

at 4th week.

Result: The age of the patients ranged from 18 to 60 years with the mean age of 32.5 (SD \pm 10.5) years. There were 52 (64.2%) male and 29 (35.8%) female with a male to female ratio was 1.8:1. The mean age of the female patients was significantly lower ($Z=1.114$; $p>0.05$) and highest 29 (35.8%) patients were in the age group of 21-30 years. Tinea corporis found in 47 (58.0%) and tinea cruris in 34 (42.0%). The mean clinical assessment score declined from 6.6 ± 1.3 at baseline; to 3.9 ± 1.2 at week 1, 0.9 ± 1.3 at week 2; 0.7 ± 1.1 at week 4. Global response was clear in 55 (67.9%), good in 25 (30.9%) and fair in 55 (67.9%). Mycological response was eradicated in 80 (98.8%) patients and persistent in only 1 (1.2%) patients. Clinical efficacy was cure in 55 (67.9%), improvement in 25 (30.9%) and failure in 1 (1.2%). Excessive pruritus occurred in 2 (2.5%) and excessive burning in 1 (1.2) after initial application of cream which gradually disappeared after subsequent application.

Biography:

Dr Sharmin Jahan has completed her MBBS at the age of 23 years from Sir Salimullah Medical College and Mitford Hospital and postgraduation study MCPS And FCPS from BCPS. She is the Consultant of Dermatology and venerology of Bangladesh Ministry of Health and family welfare. She has published 1 papers in reputed journal.