2nd International Conference on

Central Nervous System Disorders & Therapeutics

December 05-07, 2016 Dubai, UAE



Vascular factors in post stroke depression

Depression is common following an acute stroke. Post-stroke depressions (PSD) have notable impacts on the function recovery and quality of life of stroke survivors. Incidence decreased across time after stroke, but prevalence of PSD tend to be stable. Vascular factors such as diabetes, hypertension and smoking have been related to the development of PSD. Many studies have explored the association between lesion location and the incidence of PSD. For example, lesions in frontal lobe, basal ganglia and deep white matter have been related with PSD. Furthermore, cerebral microbleeds and functional changes in brain networks have also been implicated in the development of PSD. In this presentation, evidences of such association between the above structural and functional brain changes and PSD will be reviewed.

Biography

Wai Kwong Tang was appointed to Professor in the Department of Psychiatry, the Chinese University of Hong Kong in 2011. His main research areas are addictions and neuropsychiatry in stroke. He has published over 100 papers in renowned journals, and has also contributed to the peer review of 40 journals. He has secured over 20 major competitive research grants. He has served the editorial boards of five scientific journals. He was also a recipient of the Young Researcher Award in 2007, awarded by the Chinese University of Hong Kong.

tangwk@cuhk.edu.hk