

10<sup>th</sup> International Conference onNeuroscience and Neurochemistry  
&6<sup>th</sup> International Conference on Vascular Dementia February 27-March 01, 2017**A retrospective study on the demographic profile, clinical course, and management of children admitted with febrile seizures in a tertiary care hospital from 2010 to 2016****Jarren Mae R Escape**

Jarren Mae Escape, Makati Medical Center, Philippines

**Background:** Febrile seizures (FS) occur in 4-5% of children and account for the majority of seizures seen in children in emergency rooms. Local clinical practice guidelines for FS were developed in 2004. We undertook this study to look at the demographic profile of children admitted with FS, review their clinical course, diagnostic evaluations, drug management, etiology of fever, and neurological outcome. It is our hope that the information gained from this study would aid in the revision and adaptation of local clinical practice guidelines for FS.

**Objective:** To describe the clinical profile, fever etiology, clinical course, diagnostics and neurological outcome of patients admitted with febrile seizures. Data gathered was compared with clinical practice guidelines.

**Methodology:** Retrospective descriptive study that reviewed hospital records of children admitted with febrile seizures over 7 years.

**Results:** A total of 373 patients comprised the sample population. Eighty-nine percent were simple febrile seizures. Ages ranged from 3 to 91 months with the largest group in the 13-18 month old range. There was male preponderance and higher number of admissions during the rainy season. Family history was common, paternal side dominant. The most common cause of fever was upper respiratory tract infection and systemic viral illness. CBC was done in all patients. EEG's were done in 27.35% of patients; 41 % done in simple febrile seizures. Intravenous fluids and antipyretics were given and diazepam was ordered in all patients; antibiotics were given to 62.2 % of patients. Patients with complex febrile seizure are more likely to be referred to subspecialist and/or have more laboratory and imaging tests. Neurological outcome was normal.

**Conclusion:** This study showed male preponderance, increased paternal family history and seasonal variation in FS. In spite of upper respiratory tract infection and systemic viral diseases being the most common cause of fever, majority of patients received antibiotics. There was noted deviation from approved clinical practice guidelines.

**Biography**

Jarren Mae R. Escape has completed her Doctor of Medicine degree at the age of 24 years old from University of Santo Tomas. She finished her three-year residency training in Pediatrics at Makati Medical Center.

jarren\_mae@yahoo.com

**Notes:**