Cris Renata Grou Volpe et al., Prim Health Care 2017, 7:4 (Suppl)

DOI: 10.4172/2167-1079-C1-015

conferenceseries.com

4th Annual Congress & Medicare Expo on

PRIMARY HEALTHCARE AND NURSING

August 21-22, 2017 San Francisco, USA

Nursing diagnoses associated of depression indices and capacity mental in elderly people in a geriatric ambulatory clinic of Brazil

Cris Renata Grou Volpe, Marina Morato Stiva, Karolina Vicenzi Andrade, Yuri Barbalho, Luciano Ramos de Lima and Walterlânia Silva Santos University of Brasilia, Brazil

The present study aimed to identify the most prevalent nursing diagnoses in elderly patients in a geriatric outpatient clinic in the Federal district, Brazil, according to NANDA Taxonomy II relating them to the depression and mental scales. This is a descriptive, observational study of 40 elderly people over 65 years attended at the geriatric clinic. Data collection took place in March 2010 for 60 days. The scales of EDG and MEEM were used. The Pearson's chi-square was used for associations and the significance level of p<0.05. The Geriatric Depression Scale (EDG) and the Mental State Mini Exam (MMSE) were used. This study was approved by the Ethics Committee of the Health Department of the Federal District. The most prevalent nursing diagnoses were: Disturbed sensory perception; decreased cardiac output; impaired dentition; impaired memory; risk of falls and insomnia. The indexes of depressive symptoms by EDG were 35% and the cognitive alterations by the MEEM were 60%. There is a significant association between social isolation, risk of loneliness, chronic sadness and hopelessness in relation to the depression indexes demonstrated by the scale. SDs significantly associated with cognitive impairment by MMSE were poor knowledge and impaired memory. Older people tend to have cognitive deficits and depressive symptoms, especially older ones, and dependence on how much wings activities of daily living. The most prevalent related factor with impaired memory was excessive environmental changes and poor cognition was cognitive.

Biography

Cris Renata Grou Volpe is currently working as a Teacher at the University of Brasilia -UNB, Master of Health Sciences University of Sao Paulo. She has completed her PhD in Nursing at the Graduate Nursing Program (UNB). She has experience in nursing, with emphasis on fundamental nursing, medical and gerontology acting on the following topics: nursing in adult health and elderly, medicine, nursing in public health, semiotics and nursing process.

crgrou@unb.br

TI ART		4			
	O	1	Δ	0	
Τ.4	v	ı	u	Э	٠