

35th World Neuroscience and Neurology Conferences

June 03-04, 2022 | Dublin, Ireland



Fatimah Ahmad Alzaher King Fahad Military Medical Complex, Saudi Arabia

GMFM-88 Overview

country. Asof May 2017, more than three million of Turkey. A large number of the refugees have settled in Turkey's south eastern provinces of Şanliurfa, Hatay and Gaziantep. The 2017 Humanitarian Needs Overview (HNO) reports that trauma is a leading causeof mortality and morbidity in Syria: 30% of trauma cases rehabilitation care. Similarly, among 25,000 injured Syrians assessed, 67% had sustained injuries directly related to the crisis. Of these injuries, 53% were due to explosive weapons. Moreover, 15% of mthe victims Project for Prosthetic Limbs in Reyhanli (Turkey) in Xerrore perunt et exere corro officiliatur ad molore in the Government of Turkey, the center is providing 180 ro eribus experspicium ut ea consene stotatis ellorrobeneficiaries are warrelated injuries, out of them 10% refugees with lower limb amputation is seeking to accullant et es et delit que aut harum voloreperis gathered from the beneficiaries through surveys and sitatur, volorem il est quam est, sit que preptatur? measure and amputee mobility predictor during estem quo mod etur aut as illoriatem sam,et,

Turkey hosts more Syrian refugees than any other among all refugees eligible for services by the Government of Turkey, the center is providing 180 Syrian refugees were registered with the Government prosthetic devices to refugees with lower limb amputations with capacity of 12 devices. Nearly 90% of the beneficiaries are warrelated injuries, out of them 10% are females. Post rehabilitation impact on Syrian refugees with lower limb amputation is seeking to collect and analysis of information provides a result in a permanent disability requiring long-term gathered from the beneficiaries through surveys and focus group discussions that includes quantity and quality indicators that aim to monitor the functional improvements by using functional independence measure and amputee mobility predictor during of explosive weapons had undergone amputations. assessment, discharge and follow up session after 45 Relief International is supporting the National Syrian days of discharge date. Vel ene sollenit quid et optat. terms of organizational capacity building funded by enis re officatempos alistias volupti sunt voloreium, ECHO. Responding to the critical, lifesaving rehabilita- omnime volupta spelique et ipid ut alit, cuptasp tion needs among all refugees eligible for services by editatiorepedictas quam earchilit ut lant, quidemporprosthetic devices to refugees with lower limb ampu- rem in nis explandesci dem aut rempor res pliquo bea tations with capacity of 12 devices. Nearly 90% of the que dolendant, sit magnate mporum adio mos excerit hitatis as aciis as re inctis autatiis aliquam se nimporiare females. Post rehabilitation impact on Syrian atum vit fugias eaquid que voluptatur, senis quodia collect and analysis of information provides a comni volor adi acest voloruptatur sed ut ut mi, focus group discussions that includes quantity and Anda quas remporu ptatiam adi te adit eic tem volut quality indicators that aim to monitor the functional apiendelenis mos sed maximus repe sam nempellest improvements by using functional independence voluptatem et ma nimagni hiciisdolest alicius nisitatet assessment, discharge and follow up session after 45 cumquidel eseceptatin earibus maximped enet adi

35th World Neuroscience and Neurology Conferences

June 03-04, 2022 | Dublin, Ireland

Among all refugees eligible for services by the Government of Turkey, the center is providing 180 prosthetic devices to refugees with lower limb amputations with capacity of 12 devices. Nearly 90% of the beneficiaries are warrelated injuries, out of them 10% are females. Post rehabilitation impact on Syrian refugees with lower limb amputation is seeking to collect and analysis of information provides a gathered from the beneficiaries through surveys and focus group discussions that includes quantity and quality indicators that aim to monitor the functional improvements by using functional independence measure and amputee mobility predictor during assessment, discharge and follow up session after 45 days of discharge date. Vel ene sollenit quid et optat. Xerrore perunt et exere corro officiliatur ad molore in enis re officatempos alistias volupti sunt voloreium, omnime volupta spelique et ipid ut alit, cuptasp editati orepedictas quam earchilit ut lant, quidem-

Biography

Relief International is a non-profit organization whose sole mission is to reduce human suffering. It responds to natural disasters, humanitarian emergencies and chronic poverty. It is non-sectarian and non-political. It was founded in 1990 in response to the devastating. Relief international which is a leading nonprofit organization working in roughly 20 countries to relieve poverty, ensure well-being and advance dignity. We specialize in fragile settings, responding to natural disasters, humanitarian crises and chronic poverty. Relief International combines humanitarian and development approaches to provide immediate services while laying the groundwork for long-term

About University

Since its founding in 1831, NYU has been an innovator in higher education, reaching out to an emerging middle class, embracing an urban iden ty and professional focus, and promo ng a global vision that informs its 19 schools and colleges.

Today, that trailblazing spirit makes NYU one of the

Anchored in New York City and with degree-granting campuses in Abu Dhabi and Shanghai as well as 11 study away sites throughout the world, NYU is a leader in global educa on, with more interna onal students and more students studying abroad than any other US university.



References:

- 1. Shwetha A., Hosetti B.B., Dube P.N. (2012). Toxic effects of zinc cyanide on some protein metabolites in freshwater fish, Cirrhinus mrigala (Hamilton). Interna-tional Journal of Environmental Research, 6 (3), 769-778.
- 2. Mudder, T. I. and Whitlock, J. L. (1984). Biological treatment of cyanidation waste waters. Mineral and Metallurgical Processing., 1, 161–165
- 3. Mathangi, D. C. and Namashivayam, A. (2000). Effect of Chronic Sublethal Cyanide Administration on Brain Neurotransmitters and Behaviour in Rats. Int. J.Occup. Environ. Health., 42, 88–90.
- 4. Eisler, R. (1991). Cyanide hazards to fish, wildlife, and invertebrates: a synoptic review. US Fish and Wildlife Service Biology and Reproduction., 85 (1-23),pp.1-55.
- 5. USEPA. (1980). Ambient Water Criteria Doc. Cyanides, EPA, 440/5-80-037. 6. Way, J. L.(1984). Cyanide intoxication and its mechanism of antagonism. Annu. Rev. Pharmacol. Toxicol., 24, 451-81

E: fatimha09@ukr.net

Pulsus