Access

World Congress on Toxicology and Applied Pharmacology

September 05-06, 2022 | Paris, France



Fatimah Ahmad Alzaher

King Fahad Military Medical Complex, Saudi Arabia

GMFM-88 Overview

country. Asof May 2017, more than three million of Turkey. A large number of the refugees have settled in Turkey's south eastern provinces of Sanliurfa, Hatay and Gaziantep. The 2017 Humanitarian Needs Overview (HNO) reports that trauma is a leading causeof mortality and morbidity in Syria: 30% of trauma cases rehabilitation care. Similarly, among 25,000 injured Syrians assessed, 67% had sustained injuries directly related to the crisis. Of these injuries, 53% were due to explosive weapons. Moreover, 15% of mthe victims beneficiaries are warrelated injuries, out of them 10%

Turkey hosts more Syrian refugees than any other among all refugees eligible for services by the Government of Turkey, the center is providing 180 Syrian refugees wereregistered with the Government prosthetic devices to refugees with lower limb amputations with capacity of 12 devices. Nearly 90% of the beneficiaries are warrelated injuries, out of them 10% are females. Post rehabilitation impact on Syrian refugees with lower limb amputation is seeking to collect and analysis of information provides a result in a permanent disability requiring long-term gathered from the beneficiaries through surveys and focus group discussions that includes quantity and quality indicators that aim to monitor the functional improvements by using functional independence measure and amputee mobility of explosive weapons had undergone amputations. predictor during assessment, discharge and follow Relief International is supporting the National Syrian up session after 45 days of discharge date. Vel ene Project for Prosthetic Limbs in Reyhanli (Turkey) in sollenit quid et optat. Xerrore perunt et exere corro terms of organizational capacity building funded by officiliatur ad molore in enis re officatempos alistias ECHO. Responding to the critical, lifesaving rehabilita- volupti sunt voloreium, omnime volupta spelique tion needs among all refugees eligible for services by et ipid ut alit, cuptasp editati orepedictas quam the Government of Turkey, the center is providing 180 earchilit ut lant, quidempor- ro eribus experspicium prosthetic devices to refugees with lower limb ampu- ut ea consene stotatis ellorro- rem in nis explandesci tations with capacity of 12 devices. Nearly 90% of the dem aut rempor res pliquo bea que dolendant, sit magnate mporum adio mos excerit hitatis as aciis as are females. Post rehabilitation impact on Syrian re inctis autatiis aliquam se nimpori- atum vit fugias refugees with lower limb amputation is seeking to eaquid que voluptatur, senis quodia accullaut et es collect and analysis of information provides a et delit que aut harum voloreperis comni volor gathered from the beneficiaries through surveys and adi acest voloruptatur sed ut ut mi, sitatur, focus group discussions that includes quantity and volorem il est quam est, sit que preptatur? Anda quality indicators that aim to monitor the functional quas remporu ptatiam adi te adit eic tem volut improvements by using functional independence apiendelenis mos sed maximus repe sam nempellest measure and amputee mobility predictor during voluptatem et ma nimagni hiciisdolest alicius nisitatet assessment, discharge and follow up session after 45 estem quo mod etur aut as illoriatem sam, et, cumquidel eseceptatin earibus maximped enet adi

Pulsus

Volume10 Issue 1

Journal of Heart Research https://pharmacology.toxicologyconferences.com

Access

World Congress on Toxicology and Applied Pharmacology

September 05-06, 2022 | Paris, France

Among all refugees eligible for services by the Government of Turkey, the center is providing 180 prosthetic devices to refugees with lower limb amputations with capacity of 12 devices. Nearly 90% of the beneficiaries are warrelated injuries, out of them 10% are females. Post rehabilitation impact on Syrian refugees with lower limb amputation is seeking to collect and analysis of information provides a gathered from the beneficiaries through surveys and focus group discussions that includes quantity and quality indicators that aim to monitor the functional improvements by using functional independence measure and amputee mobility predictor during assessment, discharge and follow up session after 45 days of discharge date. Vel ene sollenit quid et optat. Xerrore perunt et exere corro officiliatur ad molore in enis re officatempos alistias volupti sunt voloreium, omnime volupta spelique et ipid ut alit, cuptasp editati orepedictas quam earchilit ut lant, quidem-

Biography

Relief International is a non-profit organization whose sole mission is to reduce human suffering. It responds to natural disasters, humanitarian emergencies and chronic poverty. It is non-sectarian and non-political. It was founded in 1990 in response to the devastating. Relief international which is a leading nonprofit organization working in roughly 20 countries to relieve poverty, ensure well-being and advance dignity. We specialize in fragile settings, responding to natural disasters, humanitarian crises and chronic poverty. Relief International combines humanitarian and development approaches to provide immediate services while laying the groundwork for long-term

About University

Since its founding in 1831, NYU has been an innovator in higher education, reaching out to an emerging middle class, embracing an urban iden ty and professional focus, and promo ng a global vision that informs its 19 schools and colleges.

Today, that trailblazing spirit makes NYU one of the

Anchored in New York City and with degree-granting campuses in Abu Dhabi and Shanghai as well as 11 study away sites throughout the world, NYU is a leader in global educa on, with more interna onal students and more students studying abroad than any other US university.



References:

Pulsus

1. Shwetha A., Hosetti B.B., Dube P.N. (2012). Toxic effects of zinc cyanide on some protein metabolites in freshwater fish, Cirrhinus mrigala (Hamilton). Interna-tional Journal of Environmental Research, 6 (3), 769-778.

2. Mudder, T. I. and Whitlock, J. L. (1984). Biological treatment of cyanidation waste waters. Mineral and Metallurgical Processing., 1, 161–165 3. Mathangi, D. C. and Namashivayam, A. (2000). Effect of Chronic Sublethal Cyanide Administration on Brain Neurotransmitters and Behaviour in Rats. Int. J.Occup. Environ. Health., 42, 88–90.

4. Eisler, R. (1991). Cyanide hazards to fish, wildlife, and invertebrates: a synoptic review. US Fish and Wildlife Service Biology and Reproduction., 85 (1-23), pp.1-55.

5. USEPA. (1980). Ambient Water Criteria Doc. Cyanides, EPA, 440/5- 80-037. 6. Way, J. L.(1984). Cyanide intoxication and its mechanism of antagonism. Annu. Rev. Pharmacol. Toxicol., 24, 451-81

E: fatimha09@ukr.net

Journal of Heart Research https://pharmacology.toxicologyconferences.com

Volume 10 Issue 1